

CLOVER WEED CONTROL

(Including arrowleaf, crimson, red, white, and other clovers)

Tim R. Murphy, Extension Agronomist-Weed Science

USE STAGE/ AND HERBICIDE	BROADCAST RATE/ACRE		REMARKS AND PRECAUTIONS
	AMOUNT OF FORMULATION	POUNDS ACTIVE INGREDIENT	
PREPLANT INCORPORATED			
EPTC (Eptam)	3.5 pts.	3.1	For winter annual grass and some broadleaf weed control, incorporate 2 to 3 inches deep before planting. Do not use on white Dutch clover. Do not use if a grass or grain crop is to be planted with the clover.
benefin (Balan 60DF)	2.0 lbs.	1.2	May be used on alsike, ladino, and red clover. For winter annual grass and some broadleaf weed control, incorporate 2 to 3 inches deep before planting. Do not use if a grass or grain crop is to be planted with the clover.
POSTEMERGENCE - Seedling Clovers			
imazethapyr (Pursuit) 2EC	3.0 - 6.0 fl. ozs.	0.047 - 0.094	Apply to seedling clover with a minimum of two trifoliolate leaves and when weeds are 1 to 3 inches tall or before rosette forming weeds exceed 3 inches in diameter. Pursuit requires a 1 hour rain free period. Pursuit may cause a temporary reduction in height or slight leaf yellowing. DO NOT feed, graze, or harvest clover for 30 days after application. Add a nonionic surfactant at 0.25% v/v or a crop oil concentrate at 1.25% v/v to the spray solution. If replanting is necessary in a field treated with Pursuit, do not plant alfalfa for 4 months following the application of Pursuit.
pronamide (KERB) 50W	1.0 - 1.5 lbs.	0.5 - 0.75	For preemergence control of winter annual weeds, spray prior to weed emergence in November or December. Do not graze or cut for hay for 120 days after treatment. In fall-seeded clovers, applications should be made after clover has reached the first trifoliolate leaf stage. In spring seedings, applications should be delayed until the following fall or early winter.
sethoxydim (Poast) 1.5 lb./gal. (Poast Plus) (Sethoxydim E-Pro) (Sethoxydim G-Pro) 1.0 lb./gal.	1.0 - 1.5 pts. 1.5 - 2.25 pts. 1.5 - 2.25 pts. 1.5 - 2.25 pts.	0.19 - 0.3	Apply with crop oil concentrate at 2.0 pts./acre to control annual and perennial grasses. Use the high rate for johnsongrass and bermudagrass. Do not apply to grasses growing under drought-stressed conditions. Apply to annual grasses less than 8 inches tall. Do not apply within 7 days of feeding, grazing, or harvesting undried forage, or within 20 days of cutting for dry hay.
2,4-DB (Butyrac 200)	2.3 qts.	1.0 - 1.5	For postemergence control of winter or summer annual broadleaf weeds, spray emerged weeds less than 3 inches tall. 2,4-DB will not control wild radish, chickweed, or mustards. Do not graze or cut for hay for 60 days after application. Do not apply to clovers grown for seed. Rainfall or irrigation within 7 to 10 days after treatment can cause injury. DO NOT add wetting agents or surfactants to the spray mix. Apply after clover is in the 2 to 4 trifoliolate leaf stage.
POSTEMERGENCE - Established Clovers			
imazethapyr (Pursuit) 2EC	3.0 - 6.0 fl. ozs.	0.047 - 0.094	Apply to clover with a minimum of two trifoliolate leaves and when weeds are 1 to 3 inches tall or before rosette forming weeds exceed 3 inches in diameter. Pursuit requires a 1 hour rain free period. Pursuit may cause a temporary reduction in height or slight leaf yellowing. DO NOT feed, graze, or harvest clover for 30 days after application. Add a nonionic surfactant at 0.25% v/v or a crop oil concentrate at 1.25% v/v to the spray solution. If replanting is necessary in a field treated with Pursuit, do not plant alfalfa for 4 months following the application of Pursuit.
sethoxydim (Poast) 1.5 lb./gal. (Poast Plus) (Sethoxydim E-Pro) (Sethoxydim G-Pro) 1.0 lb./gal.	1.0 - 1.5 pts. 1.5 - 2.25 pts. 1.5 - 2.25 pts. 1.5 - 2.25 pts.	0.19 - 0.3	Apply with crop oil concentrate at 2.0 pts./acre to control annual and perennial grasses. Use the high rate for johnsongrass and bermudagrass. Do not apply to grasses growing under drought-stressed conditions. Apply to annual grasses less than 8 inches tall. Do not apply within 7 days of feeding, grazing, or harvesting undried forage, or within 20 days of cutting for dry hay.

pronamide (KERB) 50W	1.0 - 1.5 lbs.	0.5 - 0.75	Controls winter annual grasses and some broadleaf weeds in clovers, birdsfoot trefoil and crown vetch. Apply from November through February. DO NOT graze or harvest for hay for 120 days after application. KERB is a restricted use herbicide.
-------------------------	----------------	------------	--